

AD-A274 651



DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

2

Information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, Send comments regarding this burden estimate or any other aspect of this collection of information, including this burden estimate, Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2. REPORT DATE January 6, 1994		3. REPORT TYPE AND DATES COVERED Progress Report: 1/1/93 - 12/31/93	
4. TITLE AND SUBTITLE Nonlinear Ocean Waves		5. FUNDING NUMBERS	
6. AUTHOR(S) Harvey Segur			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Colorado Boulder, CO 80309-0526		8. PERFORMING ORGANIZATION REPORT NUMBER 153-4647	
9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES) Department of the Navy Office of the Chief of Naval Research 800 N. Quincy Street Code 1511: KIJ Arlington, VA 22217-5000		10. SPONSORING MONITORING AGENCY REPORT NUMBER N00014-92-J-1274	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Two major papers were written during the last year, describing our work under this contract in our two main areas of research: (a) numerically induced chaos, and (b) waves in shallow water. This progress report summarizes the contents of those two papers. <div style="text-align: center;">DTIC SELECTE JAN 11 1994 S B D</div>			
14. SUBJECT TERMS		15. NUMBER OF PAGES 3	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT

Nonlinear Ocean Waves
Progress Report: 1/1/93-12/31/93

The research conducted under this contract splits naturally into two main topics: (a) numerically induced chaos; and (b) waves in shallow water. During 1993, we wrote major papers on each of these topics. This report summarizes the contents of these two papers.

(a) Numerically induced chaos

The cubic-nonlinear Schrödinger equation in one dimension,

$$i\partial_t\psi + \partial_x^2\psi + 2|\psi|^2\psi = 0, \quad (\text{NLS})$$

arises in several physical contexts, including the evolution of nearly monochromatic, one-dimensional waves in deep water. The equation is known to be a completely integrable Hamiltonian system, with infinitely many constants of the motion, *etc.* With spatially periodic boundary conditions, one can show that solutions of the equation that are quasi-periodic in time are dense among the set of all solutions, and that the system is demonstrably nonchaotic.

Even so, Ablowitz, Schober, and Herbst (1993) showed that for certain ranges of initial data, discrete numerical approximations of NLS are chaotic, even though the continuous system is nonchaotic. Moreover, they showed that the observed chaotic behaviour occurs for a variety of numerical schemes, even schemes that are themselves completely integrable, and even though these schemes preserve the constants of the motion to a very high accuracy. Even worse, tiny (10^{-16}) roundoff errors grow rapidly, and render numerical simulations meaningless within a few time units.

In other words, even though one can prove that the solution of NLS that evolves from particular initial data is well-behaved (and nonchaotic), numerical approximations of this solution are chaotic, and are completely unreliable after a few time units. We reiterate that this chaotic behaviour appears for every kind of discretization tested. The fact that it appears even for an extremely well-behaved system like NLS raises profound questions about claims of chaotic behaviour in other systems, especially when these claims are based on numerical simulations.

(b) Waves in shallow water

The objective of this work has been to develop a model of waves in shallow water that is both computationally feasible and physically realistic, without resorting to the usual restrictions to

94-01061

3R

waves of infinitesimal amplitude or to waves with one-dimensional surface patterns. The theoretical basis for all of this work is the fact that the Kadomtsev-Petviashvili equation,

$$\partial_x(\partial_t u + u\partial_x u + \partial_x^3 u) + \partial_y^2 u = 0, \quad (\text{KP})$$

describes approximately the evolution of waves of moderate amplitude as they propagate primarily in one direction in shallow water of uniform depth. In addition, the equation is known to admit large families of exact periodic and quasi-periodic solutions. The simplest of these ("genus 1") are the cnoidal waves found by Korteweg and deVries a century ago. The next level of complication ("genus 2") includes all KP solutions of permanent form. Typically, these waves of permanent form have two-dimensional surface patterns, and they are characterised by 8 real parameters.

In a paper now under review, Hammack, McCallister, Scheffner, and Segur (1993) demonstrate experimentally the existence of finite-amplitude waves of (nearly) permanent form in shallow water of uniform depth. The surface patterns of these waves are two-dimensional and spatially periodic. The basic template of a wave is hexagonal, but the wave patterns need not be symmetric about the direction of propagation, as required in previous studies. The waves are easy to generate, and they seem to be stable to perturbations. We know of no previous demonstration of genuinely two-dimensional, asymmetric waves of permanent form in shallow water. (In fact, the KP equation also arises in plasma physics and elsewhere. We know of no demonstration of the existence of two-dimensional, periodic waves of permanent form in a plasma, or in any other context in which the KP equation arises. This seems to be the first experimental demonstration of these waves in any physical context.)

In the same paper, the authors also set forth an algorithm to infer the 8 parameters of a KP solution of genus 2 from specific wave-gauge measurements, and they show that the KP solutions so obtained describe the measured waves with reasonable accuracy, even outside the putative range of validity of the KP equation.

References:

- M.J. Ablowitz, C. Schober, and B.M. Herbst, "Numerical chaos, roundoff errors, and homoclinic manifolds," *Phys. Rev. Lett.*, 71, pp. 2683-2686, 1993
- J. Hammack, D. McCallister, N. Scheffner, and H. Segur, "Two-dimensional periodic waves in shallow water, part 2: asymmetric waves," *J. Fluid Mech*, *sub judice*, 1993

For	
I	<input checked="" type="checkbox"/>
	<input type="checkbox"/>
	<input type="checkbox"/>
ion	
on/	
ity Codes	
and/or	
Special	

DTIC QUALITY INSPECTED 5

01-

A-1